

Rational coefficients of two-body energies in diagonal matrix elements of j^n configurations

Igal Talmi
Weizmann Institute of Science,
Rehovot, Israel

February 25, 2010

Abstract

Matrix elements of a two-body interaction between states of the j^n configuration (n fermions in the j -orbit) are functions of two-body energies. In many cases, diagonal matrix elements are linear combinations of two-body energies, whose coefficients are rational and positive numbers. It is shown that if in the j^n configuration there is only one state with a given spin J , its eigenvalue (the diagonal matrix element) has rational and positive coefficients of two-body energies. The situation in cases with several J -states in the j^n configuration is more complicated and is discussed in detail. States with this property are identified and constructed. In particular, the relevance to states in the seniority scheme is shown.

1 Introduction

A system of n identical fermions in a given j -orbit is considered. Such j^n configurations of protons or neutrons occur, for instance, in the shell model. Any

configuration contains a certain number of fully antisymmetric independent states allowed by the Pauli principle. We consider shell model Hamiltonians which contain mutual interactions between the particles. The interactions considered are rotationally invariant and hence, eigenstates of the Hamiltonian have definite values of angular momentum J . The mutual interaction is considered to be a perturbation. Hence, only the sub-matrix of the Hamiltonian defined by the states of the j^n configuration is taken into account. The states considered here are the states which define this sub-matrix.

If there is only one state with given J in the j^n configuration, it is an eigenstate of the sub-matrix of the Hamiltonian. Its eigenvalue is given by the corresponding diagonal element of the sub-matrix. If there are several independent states with the same value of J , they define a smaller sub-matrix whose diagonalization yields the eigenstates for a given mutual interaction. If two-body mutual interactions are considered, elements of the Hamiltonian sub-matrix of the j^n configuration are functions of two-body matrix elements. Due to the rotational invariance of the interaction, the only non-vanishing two-body matrix elements are between two-body states with the same value of J . Thus, these two-body matrix elements are the energies of the two-body configuration.

In a recent paper [1], Zamick and Van Isacker consider partial dynamical symmetries of certain states of the $(9/2)^4$ configuration. They used their results to calculate also the energies of the $J = 2$ and $J = 4$ states of the $(7/2)^4$ configuration. They express these energies as linear combinations of two-body energies and in their arXiv: 0803.1569v1 (nucl-th) version they "note that this derivation constitutes a proof that the coefficients...in the energy expression...must be rational numbers".

This feature is quite general. There are many cases where diagonal matrix element in j^n configurations are linear combinations of two-body energies whose coefficients are rational and positive numbers. This feature has little relevance to numerical calculations but it still has theoretical interest. The

aim of this paper is to study this property of states, to find the cases in which diagonal elements are linear combinations of two-body energies whose coefficients are positive rational numbers.

There is a simple way to calculate the diagonal matrix elements of a two-body interaction in a system of n fermions in a j -orbit. It is based on the use of m -scheme wave functions. These are determinantal wave functions characterized by a set of quantum numbers $m_1, m_2, \dots, m_n$, which are all different. These wave functions are discussed in great detail in ref.[2] where matrix elements of a two body interaction between such states are derived. It was shown there that diagonal matrix elements are equal to the $\sum \langle m_i m_k | V | m_i m_k \rangle$ where the summation is over all pairs of m -values in the state considered.

Matrix elements of a rotationally invariant two-body interaction between m -states with given total $M = \sum m_i$ form a sub-matrix. Its trace is equal to the sum of diagonal elements taken between all states with given M . States of the j^n configuration with given values of J and M , $J \geq M$, are linear combinations of these m -states. The linear transformation, from m -states to states with definite J and M , is unitary or rather orthogonal. Hence, the trace of the sub-matrix is invariant and may be used to calculate diagonal matrix elements of a rotationally invariant, M -independent interaction in the J -scheme.

Subtracting the trace of the $M = J - 1$ sub-matrix from the one of $M = J$, yields the sum $\sum \langle j^n a J, M | V | j^n a J, M \rangle$ where a characterizes the state if there are several states with given J in the j^n configuration. If there is only one such state, this procedure yields its interaction energy. Otherwise, only the sum of energies of all states with given J is obtained. This case will be discussed in detail in the following. The diagonal matrix elements in the m -scheme may be expressed in terms of matrix elements of two particles coupled to a given J value. The two-particle matrix elements in the m -scheme may be transformed by using Clebsch-Gordan coefficients

as follows.

$$\begin{aligned} & \langle m_i m_k | V | m_i m_k \rangle = \\ & \sum \langle j, m_i, j, m_k | j, j, J, M = m_i + m_k \rangle^2 \langle j^2 JM | V | j^2 JM \rangle \end{aligned} \quad (1)$$

Since the m -states in (1) are antisymmetric, the sum on the r.h.s. of (1) is only over *even* values of J (states of two fermions with *odd* J values are symmetric). Thus, the coefficients of the $V(J) = \langle j^2 JM | V | j^2 JM \rangle$ in the diagonal elements in the m -scheme are equal to sums of squares of Clebsch-Gordan coefficients. From formulae for these coefficients (e.g. (15.36), ref.[2]) it is seen that their squares are rational functions of their arguments. These arguments are the 3 J -values and 3 m -values.

Thus, the sum $\sum_J \langle j^n aJ, M | V | j^n aJ, M \rangle$ is a linear combination of two-body matrix elements $V(J) = \langle j^2 JM | V | j^2 JM \rangle$ which are independent of M , whose coefficients are rational and positive numbers. If there is only one state with given J in the j^n configuration, it has the "rational property" considered here. Its eigenvalue of a two-body interaction is a linear combination of the $V(J)$ with positive rational coefficients. If there are several such states, it was shown above that this property holds only for the sum of their diagonal elements. It will be shown in this paper which of these states has this property. It will also be shown below, that in such cases, not any arbitrary state may have this property.

States in the j^n configuration with given J must be considered individually. This may be achieved by using coefficients of fractional parentage (c.f.p.), introduced by Bacher and Goudsmit [3] and intensively applied to atomic spectra by Racah[4]. An antisymmetric wave function with n particles and given J may be expanded as

$$\psi(j^n aJ) = \sum [j^{n-1}(bJ_1)jJ] \psi(j^n aJ) \psi(j^{n-1}(bJ_1)j_n aJ) \quad (2)$$

In (2), a and b are used to specify uniquely J and J_1 states if there are several of them. The wave functions on the r.h.s. of (2) are fully antisymmetric functions of particles 1 to $n-1$. They are usually not antisymmetric in all n particles but their linear combination in (2) is.

Using the expansion (2), matrix elements in states of the j^n configuration may be expressed in terms of those in the j^{n-1} configuration. Due to the full antisymmetry of (2), it is possible to calculate matrix elements of ΣV_{ik} for $i, k < n$ whose number is $(n-1)(n-2)/2$ and multiply the result by $n(n-1)/(n-1)(n-2)=n/(n-2)$ to obtain the full value of the matrix element. This leads to

$$\begin{aligned} & \langle j^n a_n J | \Sigma V_{ik} | j^n b_n J \rangle = \\ & n/(n-2) \sum_{a_{n-1} J_1} [j^{n-1}(a_{n-1} J_1) j J] \{ j^n a_n J \} [j^{n-1}(b_{n-1} J_1) j J] \{ j^n b_n J \} \times \\ & \langle j^{n-1} a_{n-1} J_1 | \Sigma V_{ik} | j^{n-1} b_{n-1} J_1 \rangle \end{aligned} \quad (3)$$

Using c.f.p. the matrix elements in (3) may be further reduced to those in the j^{n-2} configuration and so on until the j^2 configuration is reached. A direct result of (3) follows if there is only one state with given J in the configurations involved. Then, diagonal matrix elements have the property considered here if the squares of the c.f.p. are rational numbers. Since this was proved above, the case with several states with the same value of J should be considered. This will be carried out in the following, starting with the simple case of $n=3$ and proceeding to higher n values.

2 A simple case: the j^3 configuration

A state with given J may be obtained by starting from a *principal parent* with given J_0 in the j^2 configuration, coupling to it j_3 and antisymmetrizing the function $\psi(j^2(J_0)j_3J)$. If the resulting state is allowed, the wave function does not vanish and its c.f.p. are given by (e.g. (26.11), ref.[2] or (15.10), ref.[5])

$$\begin{aligned} & [j^2(J_1)jJ] \{ j^3[J_0]J \} = \\ & N(J_0, J) [\delta(J_1, J_0) + 2[(2J_0 + 1)(2J_1 + 1)]^{1/2} \{ jjJ_1/JjJ_0 \}] \end{aligned} \quad (4)$$

where the normalization factor is given by

$$N(J_0, J) = [3 + 6(2J_0 + 1) \{ jjJ_0/JjJ_0 \}]^{-1/2} \quad (5)$$

and $\{abc/def\}$ are $6j$ -symbols (Racah coefficients). The c.f.p. of the principal parent is obtained by putting $J_1 = J_0$ in (4). Then from (5) follows the special $n = 3$ case of the identity proved in ref.[2],

$$nN(J_0, J)[j^2(J_0)jJ|]j^3[J_0]J = 1 \quad (6)$$

To find the properties of the c.f.p., it is worth while to look at the form of the $6j$ -symbols. When Racah introduced these symbols [6] (in a slightly different form and symmetry properties), he also derived a closed formula of them in terms of their 6 arguments. In (15.38),ref.[2], it is expressed as

$$\{abc/def\} = \Delta(abc)\Delta(aef)\Delta(dbf)\Delta(dec)R(abcdef) \quad (7)$$

where

$$\Delta(abc) = [(a+b-c)!(b+c-a)!(c+a-b)!/(a+b+c+1)!]^{1/2} \quad (8)$$

and R is a rational function of the 6 arguments. Hence, the square of a $6j$ -symbol is a rational function of its arguments. If a state obtained by antisymmetrization is not allowed by the Pauli principle, the wave function vanishes. This always happens for states with $J=3j-4$ or $J=\frac{1}{2}$. The unnormalized c.f.p. all vanish due to the properties of the rational function in(7). From (7) and (8) follow some simple properties of $6j$ -symbols, like

$$\{jjJ_0/JjJ_0\} = \Delta(jjJ_0)^2\Delta(JjJ_0)^2R(jjJ_0JjJ_0) \quad (9)$$

From (9) follows that the c.f.p. of the principal parent, with $J_1 = J_0$, is a square root of a rational function of j , J_0 and J . This is true also for the normalization coefficient (5). Therefore, also the other c.f.p., with $J_1 \neq J_0$, share this feature since their squares are equal to

$$4N(J_0, J)^2(2J_0+1)(2J_1+1)\{j, j, J_1/J, j, J_0\}^2 \quad (10)$$

Thus, the squares of the c.f.p. of a state obtained by antisymmetrization of a principal parent are rational functions of the angular momenta. Hence, due to (3), such states have the property considered here. Their eigenvalues of a

two-body interaction are linear combinations of the $V(J) = \langle j^2 J | V | j^2 J \rangle$ with coefficients which are positive rational numbers. For $n=3$, the relation (3) yields directly these linear combinations as follows

$$\begin{aligned} & \langle j^3 a J | \sum V_{ik} | j^3 a J \rangle = \\ & 3 \sum [j^2(J_1) j J] \{ j^3 a J \} [j^2(J_1) j J] \{ j^3 a J \} \langle j^2 J_1 | V | j^2 J_1 \rangle \end{aligned} \quad (11)$$

If there is only one state with given J in the j^3 configuration, the proof that it has the property considered, is complete as already shown in the preceding section. In that case, starting with all possible principal parents, those with J_0 which can yield J when coupled with j , the same state is obtained after antisymmetrization (with a possible change of overall sign). If there are several states with given J in the j^3 configuration, they may be constructed by antisymmetrizing *several* principal parents. The diagonal matrix element of each of them is given by the relation (11). Still, the states obtained in this way need not be orthogonal. It will be shown now that states obtained in the orthogonalization procedure have also the property considered.

The orthogonalization of states with the same value of J constructed from principal parents is simplified by the following property of such c.f.p. Due to (4), they may be expressed, for $J_1 \neq J_0$, as

$$\begin{aligned} & [j^2(J_1) j J] \{ j^3 [J_0] J \} = \\ & N(J_0, J) 2[(2J_0 + 1)(2J_1 + 1)]^{1/2} \{ j j J_1 / J j J_0 \} = \\ & N(J_0) (2J_0 + 1)^{1/2} \Delta(j j J_0) \Delta(J j J_0) (2J_1 + 1)^{1/2} \Delta(j j J_1) \Delta(J j J_1) R \end{aligned} \quad (12)$$

where $N(J_0) = N(J_0, J)$ (also in the following) and $R = R(j j J_1 J j J_0)$. Thus, these c.f.p. are products of a rational function which multiplies a term which depends only on J_0 and another one which depends only on J_1 . The latter term is the same for all states with various values of J_0 .

To make a state $|b\rangle$ orthogonal to a given state $|a\rangle$, the latter multiplied

by $\langle a|b \rangle$ should be subtracted from the former. The orthogonalized state

$$|b' \rangle = |b \rangle - \langle a|b \rangle |a \rangle \quad (13)$$

should then be normalized. Using c.f.p., the scalar product of the two states is given by

$$\langle j^n a J | j^n b J \rangle = \Sigma_{a_1 J_1} [j^{n-1}(a_1 J_1) j J] [j^n a J] [j^{n-1}(a_1 J_1) j J] [j^n b J] \quad (14)$$

The state $|a \rangle$ is taken to be constructed from the principal parent $\psi(j^2(J_0)j_3 J)$ and the $|b \rangle$ state from $\psi(j^2(J'_0)j_3 J)$. Making use of the c.f.p. (4) we find that every term in the summation in (14), for $J_1 \neq J_0, J_1 \neq J'_0$ is given by

$$\begin{aligned} & N(J_0)N(J'_0)2[(2J_0+1)(2J_1+1)]^{1/2}\{jjJ_1/JjJ_0\} \times \\ & 2[(2J'_0+1)(2J_1+1)]^{1/2}\{jjJ_1/JjJ'_0\} \end{aligned} \quad (15)$$

From the properties (7) of $6j$ -symbols follows that the product of the two of them in (15) can be expressed as

$$\begin{aligned} & \{jjJ_1/JjJ_0\}\{jjJ_1/JjJ'_0\} = \Delta(jjJ_1)\Delta(jjJ_0)\Delta(JjJ_1)\Delta(JjJ_0) \times \\ & \Delta(jjJ_1)\Delta(jjJ'_0)\Delta(JjJ_1)\Delta(JjJ'_0)R(jjJ_1JjJ_0)R(jjJ_1JjJ'_0) = \\ & \Delta(jjJ_0)\Delta(JjJ_0)\Delta(jjJ'_0)\Delta(JjJ'_0)\Delta(jjJ_1)^2\Delta(JjJ_1)^2 \times \\ & R(jjJ_1JjJ_0)R(jjJ_1JjJ'_0) \end{aligned} \quad (16)$$

Due to the form (7) of $6j$ -symbols, the r.h.s. of (16) may be expressed as

$$\begin{aligned} & (jjJ_0/JjJ'_0/R(jjJ_0JjJ'_0))\Delta(jjJ_1)^2\Delta(JjJ_1)^2 \times \\ & R(jjJ_1JjJ_0)R(jjJ_1JjJ'_0) \end{aligned} \quad (17)$$

This expression is a $6j$ -symbol multiplied by a rational function, $R(J_1)$. Thus, every term in (14) becomes equal to

$$4N(J_0)N(J'_0)[(2J_0+1)(2J'_0+1)]^{1/2}\{jjJ_0/jjJ'_0\} \times (2J_1+1)R(J_1) \quad (18)$$

The summation of terms with values of J_1 different from J'_0 and from J_0 may now be carried out and yields a rational function multiplying

$$N(J_0)N(J'_0)[(2J_0+1)(2J'_0+1)]^{1/2}\{jjJ_0/jjJ'_0\} \quad (19)$$

The term in (14) with $J_1 = J_0$ is equal to

$$\begin{aligned} & N(J_0)N(J'_0)[1 + 2[(2J_0 + 1)(2J_0 + 1)]^{1/2}\{jjJ_0/JjJ_0\}] \times \\ & 2[(2J'_0 + 1)(2J_0 + 1)]^{1/2}\{jjJ_0/JjJ'_0\} \end{aligned} \quad (20)$$

Due to the properties of $6j$ -symbols (7), this expression is equal to a rational function multiplying the expression (19). Also the term with $J_1 = J'_0$ is equal to (19) multiplied by a (different) rational function. Thus, the summation in (14) yields a rational function multiplied by the factor (19).

To obtain the state which is orthogonal to the one characterized by J'_0 , eq.(13) is used. The unnormalized c.f.p. of that state are defined by

$$\begin{aligned} & N(J'_0)[\delta(J_1, J'_0) + 2[(2J'_0 + 1)(2J_1 + 1)]^{1/2}\{jjJ_1/JjJ'_0\}] - \\ & N(J_0)N(J'_0)[(2J_0 + 1)(2J'_0 + 1)]^{1/2}\{jjJ_0/jjJ'_0\} \times \\ & N(J_0)[\delta(J_1, J_0) + 2[(2J_0 + 1)(2J_1 + 1)]^{1/2}\{jjJ_1/JjJ_0\}]R \end{aligned} \quad (21)$$

where R is a rational function. In the case that J_1 is different from J_0 and J'_0 , (21) assumes the form

$$\begin{aligned} & N(J'_0)2[(2J'_0 + 1)(2J_1 + 1)]^{1/2}\{jjJ_1/JjJ'_0\} - \\ & N(J_0)N(J'_0)[(2J_0 + 1)(2J'_0 + 1)]^{1/2}\{jjJ_0/jjJ'_0\} \times \\ & N(J_0)2[(2J_0 + 1)(2J_1 + 1)]^{1/2}\{jjJ_1/JjJ_0\}R \end{aligned} \quad (22)$$

In (22) there are several square roots which appear twice. The product of two $6j$ -symbols is equal, according to (16) and (17) to $\{jjJ_1/jjJ'_0\}$ multiplied by a rational function. Hence, the unnormalized c.f.p. of the orthogonalized state, given by (22) are equal to the original c.f.p. multiplied by rational functions.

In the case $J_1 = J_0$, that c.f.p. is equal to

$$\begin{aligned} & N(J'_0)2[(2J'_0 + 1)(2J_0 + 1)]^{1/2}\{jjJ_0/JjJ'_0\} - \\ & N(J_0)N(J'_0)[(2J_0 + 1)(2J'_0 + 1)]^{1/2}\{jjJ_0/jjJ'_0\} \times \\ & N(J_0)[1 + 2[(2J_0 + 1)(2J_0 + 1)]^{1/2}\{jjJ_0/JjJ_0\}]R \end{aligned} \quad (23)$$

In (23), $N(J_0)^2[1 + 2[(2J_0 + 1)(2J_0 + 1)]^{1/2}\{jjJ_0/JjJ_0\}]$ is a rational function and hence, (23) is equal to the original c.f.p. multiplied by a rational function, as are the other c.f.p. Also in the case $J_1 = J'_0$, the situation is similar. In that case, (21) is equal to

$$\begin{aligned} & N(J'_0)[1 + 2[(2J'_0 + 1)(2J'_0 + 1)]^{1/2}\{jjJ'_0/JjJ'_0\}] - \\ & N(J_0)N(J'_0)[(2J_0 + 1)(2J'_0 + 1)]^{1/2}\{jjJ_0/jjJ'_0\} \times \\ & N(J_0)2[(2J_0 + 1)(2J'_0 + 1)]^{1/2}\{jjJ'_0/JjJ_0\}R \end{aligned} \quad (24)$$

The c.f.p. of the principal parent of the original state is equal to $N(J'_0)$ multiplied by a rational function. The term subtracted from it is equal to $N(J'_0)$ multiplied by the product of the rational function R and the square of a product of square roots. Hence, all c.f.p. of the unnormalized orthogonalized state are equal to the original ones multiplied by various rational functions. The resulting state should be normalized. It should be divided by the square root of the sum of squares of these c.f.p. Since the squares of the latter are rational, the normalization coefficient is the square root of a rational function. Thus, the squares of the c.f.p. of the normalized and orthogonalized state are rational and positive.

The c.f.p. of the orthogonalized and normalized states are equal, apart from a common normalization factor, to the original c.f.p. multiplied by rational functions. It may happen that certain c.f.p. vanish after this procedure. For instance, starting from a certain J_0 , in the state with J'_0 which was made orthogonal to it, the c.f.p. for $J_1 = J_0$ must vanish. If further orthogonalizations are necessary, due to existence of more independent states with the same total J , It is possible to repeat the steps taken above.

Looking at the derivations above, it is clear that not all linear combinations of states obtained from principal parents have the property considered here. The c.f.p. may become sums of square roots of relatively prime numbers and their squares will then not be rational numbers. In fact, there are states whose diagonal matrix elements are not linear combinations of two-body energies. This may happen to eigenstates obtained by diagonalization

of the submatrix, with given J , of a two-body interaction. An example is shown at the end of this section. Naturally, it is possible to construct linear combinations, other than those discussed above, which have the "rational property".

Non-diagonal matrix elements have also a simple form when taken between states sharing the property discussed here. We consider the matrix element (3) for $n=3$ taken between different states. We may take them to be the state obtained by antisymmetrizing the principal parent with J_0 and the one obtained from the J'_0 principal parent, orthogonalized as above. To calculate the non-diagonal matrix element it is possible to make use of the calculation of the overlap of the two states above. We start from the formula (3) for the special case of $n=3$

$$\begin{aligned} & \langle j^3[J_0]J | \Sigma V_{ik} | j^3[J'_0]J \rangle = \\ & 3\Sigma[j^2(J_1)jJ] \{j^3[J_0]J\} [j^2(J_1)jJ] \{j^3[J'_0]J\} \langle j^2J_1 | V | j^2J_1 \rangle \end{aligned} \quad (25)$$

Every term in the summation in (25) is equal to a corresponding terms in (14) multiplied by $\langle j^2J_1 | V | j^2J_1 \rangle$. Every term (14) is transformed into (19), which is

$$N(J_0)N(J'_0)[(2J_0+1)(2J'_0+1)]^{1/2} \{jjJ_0/jjJ'_0\} \quad (26)$$

multiplied by a rational function. Hence, the coefficient of $\langle j^2J_1 | V | j^2J_1 \rangle$ in (25) is (26) multiplied by a rational function. Thus, the non-diagonal matrix elements in the scheme of states described above, are linear combinations of two-body energies with rational functions multiplied by a common square root of a rational function.

Before proceeding to higher values of n , it is worth-while to look at some simple examples in j^3 configurations. States with $J=j$ (for n odd) and seniorities $v=1$ of the j^n configuration are obtained from principal parents with $J_0 = 0$ and $v_0 = 0$ in the j^{n-1} configuration. The seniority scheme is explained in detail in refs.[2] and [5]. It is also reviewed in the last section of this paper. In the case of $n=3$, putting $J_0 = 0$ in (4) the following results

were obtained

$$\begin{aligned}
[j^2(0)jJ = j|]j^3v = 1J = j] &= \{1 + 2\{jj0/jj0\}N(0) = \\
&[(2j - 1)/3(2j + 1)]^{1/2} \\
[j^2(J_1)jJ = j|]j^3v = 1J = j] &= \\
2(2J_1 + 1)^{1/2}\{jjJ_1/jj0\}N(0) &= -2[(2J_1 + 1)/3(2j - 1)(2j + 1)]^{1/2} \quad (27)
\end{aligned}$$

As long as $j < 9/2$, there is only one state with $J=j$ in the j^3 configuration and that state has c.f.p. given above. For higher values of j there are other states with $J=j$ and they all have seniorities $v=3$. Their number is $[(2j-3)/6]$ where $[x]$ is the largest integer not exceeding x . They may be constructed from principal parents with even values of $J_0 > 0$ and then orthogonalized to the $v=1$ state and to each other. The first part of this procedure, for any j , is carried out in ref.[5], Section 20, pages 396 - 398.

The two states with $J=9/2$ in the j^3 configuration offer an opportunity to demonstrate a statement made above. Not all states have the property considered in this paper. There are states whose diagonal matrix elements are not equal to linear combinations of two-body energies. It was mentioned that such cases occur in diagonalization of the Hamiltonian sub-matrix defined in the j^n configuration by the states with a given value of J .

Consider the 2x2 sub-matrix defined by the $v=1$ and $v=3$ states with $J=9/2$. Both diagonal matrix elements V_1 and V_3 are linear combinations of the $V(J)$ with positive rational coefficients. The non-diagonal matrix element V is also a linear combination of the $V(J)$ with rational coefficients multiplied by the square root of a rational function. The two eigenvalues with $J=9/2$ are the roots of the quadratic secular equation

$$x^2 - (V_1 + V_3)x + V_1V_3 - V^2 = 0 \quad (28)$$

They are equal to

$$\begin{aligned}
x &= \frac{1}{2}\{V_1 + V_3 \pm [(V_1 + V_3)^2 - 4V_1V_3 + 4V^2]^{1/2}\} = \\
&\frac{1}{2}\{V_1 + V_3 \pm [(V_1 - V_3)^2 + 4V^2]^{1/2}\} \quad (29)
\end{aligned}$$

Both eigenvalues are sums of linear combinations with rational coefficients and square roots of quadratic functions of the $V(J)$.

In some special cases, the square root may turn out to be a linear function of the $V(J)$. In the case of the $J = 9/2$ described above, this cannot happen. The $v = 3$ state is orthogonal to the $v = 1$ state whose principal parent is the $j^2(J_1 = 0)$ state. Hence, the c.f.p. with $J_1=0$ of the $v=3$ state vanishes. Thus, in the linear combination V_3 the coefficient of $V(0)$ vanishes. This coefficient vanishes also in the non-diagonal matrix element V . The term $V_1 - V_3$ may be expressed as a linear combination of $V(0)$ and terms with $V(J), J > 0$ even, $c_0V(0) + \Sigma c_J V(J)$. Its square is equal to $c_0^2V(0)^2 + 2c_0V(0)\Sigma c_J V(J) + (\Sigma c_J V(J))^2$. The term with V^2 has the form $(\Sigma d_J V(J))^2$ with no $V(0)$ terms, and adding it cannot change the quadratic and linear terms in $V(0)$. If V does not vanish, adding $4V^2$ to $(V_1 - V_3)^2$ cannot yield a square of a rational function of the $V(J)$. Hence, in the example considered here, the eigenvalues of a two-body interaction are definitely not linear combinations of the two-body energies.

There is no contradiction between this result and (11). There, the diagonal matrix element of a state is a linear combination of the $V(J)$ with coefficients which are squares of the c.f.p.(multiplied by 3). The states constructed in (4), as well as all states constructed in this paper, are states which form a basis of all states with given J . Any antisymmetric state may be expanded in terms of c.f.p. as in (2). The states constructed in (4), as well as others, are based on the use of principal parents. Hence, the c.f.p. are functions of the angular momenta and are independent of the two-body interaction. If there is only one state with given J , it is an eigenstate and its eigenvalue is a linear combination of the $V(J)$. If there are several such states and the Hamiltonian sub-matrix has non-vanishing non-diagonal elements, this is no longer the case. Eigenstates may be expanded in terms of c.f.p. but the latter may well be (irrational) functions of the two-body matrix elements. This is indeed the case in the example shown above.

3 General j^n configurations

In the preceding section it was shown that for the $n=3$ case, states constructed by antisymmetrizing principal parents have the property considered in this paper. Diagonal matrix elements of a two-body interaction are linear combinations of two-body matrix elements (energies) $V(J)$ whose coefficients are rational numbers. This property is still shared by such states which were orthogonalized to each other. These features follow from demonstrating that for $n=3$, the c.f.p. of these states are square roots of rational functions of the angular momenta involved. It was shown that the normalization factor is a square root of a rational function. That the c.f.p. of the principal parent is equal to the normalization factor multiplied by a rational function. There is a general relation between this c.f.p. and the normalization factor. It was proved in ref.[2], (26.23), that

$$n[j^{n-1}(a, J_0)jJ]\}j^n[a, J_0]J]N(J_0, J) = 1 \quad (30)$$

In the following it will be shown by induction that these features are shared also by c.f.p. of states in j^n configurations for $n > 3$. Starting from $n=3$ it will be shown that if these features hold for c.f.p. in the j^{n-1} configuration, they hold also for c.f.p. in the j^n configuration. The property considered in this paper, that the expectation value of a two-body interaction is then a linear combination of two-body energies whose coefficients are rational functions then follows as will be shown below.

Consider a state of the j^n configuration obtained by antisymmetrizing a principal parent $\psi(j^{n-1}(aJ_0)j_nJ)$. If the result does not vanish, its c.f.p. can be expressed by c.f.p. of states in the j^{n-1} configuration. All states considered in the following are taken to be allowed states whose wave functions do

not vanish. This relation (ref.[2] eq.(26.30) or (15.29) of ref.[5]) is

$$\begin{aligned}
& n[j^{n-1}(a_{n-1,0} J_{n-1,0})jJ_n|]j^n[a_{n-1,0} J_{n-1,0}]J_n] \times \\
& [j^{n-1}(a_{n-1,1} J_{n-1,1})jJ_n|]j^n[a_{n-1,0} J_{n-1,0}]J_n] = \\
& \delta(a_{n-1,1}, a_{n-1,0})\delta(J_{n-1,1}, J_{n-1,0}) + \\
& (n-1)\Sigma(-1)^{J_{n-1,0}} + J_{n-1,1}[(2J_{n-1,0} + 1)(2J_{n-1,1} + 1)]^{1/2} \\
& \{J_{n-2}jJ_{n-1,1}/J_njJ_{n-1,0}\}[j^{n-2}(a_{n-2}J_{n-2})jJ_{n-1,0}|]j^{n-1}a_{n-1,0}J_{n-1,0}] \times \\
& [j^{n-2}(a_{n-2}J_{n-2})jJ_{n-1,1}|]j^{n-1}a_{n-1,1}J_{n-1,1}] \quad (31)
\end{aligned}$$

The first step in calculating the c.f.p. is to put in (31), $a_{n-1,1} = a_{n-1,0}$ and $J_{n-1,1} = J_{n-1,0}$. This leads to the result

$$\begin{aligned}
& n[j^{n-1}(a_{n-1,0}J_{n-1,0})jJ_n|]j^n[a_{n-1,0}J_{n-1,0}]J_n]^2 = \\
& 1 + (n-1)(-1)^{2J_0}\Sigma(2J_{0+1})\{J_2jJ_0/JjJ_0\} \times \\
& [j^{n-2}(a_{n-2}J_{n-2})jJ_{n-1,0}|]j^{n-1}a_{n-1,0}J_{n-1,0}]^2 \quad (32)
\end{aligned}$$

As in (9), the 6 j -symbol in (32) is a rational function of its arguments. If the squares of the $n-2 \rightarrow n-1$ c.f.p. are rational, as assumed for the induction, then the c.f.p. on the l.h.s. of (32) is indeed the square root of a rational function for any n . According to (30), the normalization coefficient is also a square root of a rational function.

The c.f.p. of all states are given by the r.h.s. of (31) divided by n multiplying the c.f.p. of the principal parent. Due to (30), this amounts to multiplying the r.h.s. of (31) by the normalization coefficient $N(J_{n-1,0}J_n)$. Starting from the known c.f.p. for the $n=3$ case, it seems that in general the c.f.p. $[j^{n-1}(a_{n-1,1}J_{n-1,1})jJ_n|]j^n[a_{n-1,0}J_{n-1,0}]J_n]$ is equal to a rational function of the various angular momenta multiplied by

$$\begin{aligned}
& N(J_{n-1,0}J_n)\Pi N(J_{r0}J_{r+1,0})\Pi N(J_{r1}J_{r+1,1})\Pi[(2J_{r0} + 1)(2J_{r1} + 1)]^{1/2} \times \\
& \Pi\Delta(J_{r0}jJ_{r+1,0})\Pi\Delta(J_{r1}jJ_{r+1,1})\Delta(J_{n-1,0}jJ_n)\Delta(J_{n-1,1}jJ_n) \quad (33)
\end{aligned}$$

In the products in (33), $J_{1,0} = J_{1,1} = j$ and r goes up to $n-2$. In the following, this form of c.f.p. will be proved by induction. Apart from the first factor, (33) is a product of two expressions which are the same functions of the various J_{r0} and the various J_{r1} . The former define the state considered and the

latter, the component whose c.f.p. is selected. This feature was noticed in the $n=3$ case and is in agreement with the property of c.f.p. given by (26.21) of ref[2]. The various angular momenta belong to wave functions which were obtained by antisymmetrization of principal parents and by orthogonalization if necessary. The intermediate angular momenta supply the additional quantum numbers, denoted above by a_{r0} and a_{r1} , needed to specify the states considered. This will be the case for all states considered below.

It is worth-while to point out that the form (33) is the correct one also for the c.f.p. of the principal parent. Putting $J_{r1} = J_{r0}$ for all r , the result is a rational function multiplying the normalization factor

$$N(J_{n-1,0}J_n) = nN(J_{n-1,0}J_n)^2/nN(J_{n-1,0}J_n)$$

According to (30) this is equal to a rational function multiplying the c.f.p. of the principal parent. Hence, no special care should be paid to that c.f.p. The fact that (33) is the correct expression of all c.f.p. will be proved by induction with respect to n . Before doing it, let us draw some conclusions from the structure (33) of c.f.p.

The first conclusion which follows from (33) is that the squares of these c.f.p. are positive rational functions of the various angular momenta. This was shown above for the c.f.p. of the principal parent and this feature is shared by all of them. The aim of this paper is to show for which states the expectation values of a two-body interaction are linear combinations of two-body energies with rational coefficients. When (33) will be shown to be correct, the answer to this question will be given. States with this property are those obtained by consecutive constructions from principal parents and antisymmetrizations.

Another property of c.f.p. which follows from (33) is important for the construction of orthogonal sets of states with the same total spin J . As in the case of $n=3$, we construct such a set by choosing one state due to a principal

parent. If there are other such independent states, obtained from other principal parents, they are made orthogonal to it. The orthogonalization is carried out by using (13). First, the scalar product (14) is calculated, for the state (33) defined by J'_{r0} and the one defined by J_{r0} . It is a linear combination and each of its terms is a rational function multiplied by

$$\begin{aligned}
& N(J_{n-1,0} J_n) \Pi N(J_{r0} J_{r+1,0}) \Pi N(J_{r1} J_{r+1,1}) \Pi [(2J_{r0} + 1)(2J_{r1} + 1)]^{1/2} \times \\
& \Pi \Delta(J_{r0} j J_{r+1,0}) \Pi \Delta(J_{r1} j J_{r+1,1}) \Delta(J_{n-1,0} j J_n) \Delta(J_{n-1,1} j J_n) \times \\
& N(J'_{n-1,0} J_n) \Pi N(J'_{r0} J'_{r+1,0}) \Pi N(J_{r1} J_{r+1,1}) \Pi [(2J'_{r0} + 1)(2J_{r1} + 1)]^{1/2} \times \\
& \Pi \Delta(J'_{r0} j J'_{r+1,0}) \Pi \Delta(J_{r1} j J_{r+1,1}) \Delta(J'_{n-1,0} j J_n) \Delta(J_{n-1,1} j J_n)
\end{aligned} \tag{34}$$

In (34) all terms with J_{r1} appear twice and thus contribute only to the rational function multiplying it. Hence, the summation over $J_{n-1,1}$ in (14) may be carried out yielding for the scalar product a rational function multiplying

$$\begin{aligned}
& N(J_{n-1,0} J_n) N(J'_{n-1,0} J_n) \Pi N(J_{r0} J_{r+1,0}) \Pi N(J'_{r0} J'_{r+1,0}) \times \\
& \Pi [(2J'_{r0} + 1)(2J_{r0} + 1)]^{1/2} \Pi \Delta(J_{r0} j J_{r+1,0}) \Pi \Delta(J'_{r0} j J'_{r+1,0}) \times \\
& \Delta(J_{n-1,0} j J_n) \Delta(J'_{n-1,0} j J_n)
\end{aligned} \tag{35}$$

According to (13), the state with J_{r0} should be multiplied by the scalar product (35) and subtracted from the state with J'_{r0} which is to be orthogonalized to the other state. The factor (35) multiplied by the rational function is independent of $J_{n-1,1}$. Multiplying it by a c.f.p. of the state with J_{r0} , we obtain according to (35) and (33), a rational function multiplying

$$\begin{aligned}
& N(J_{n-1,0} J_n) N(J'_{n-1,0} J_n) \Pi N(J_{r0} J_{r+1,0}) \Pi N(J'_{r0} J'_{r+1,0}) \times \\
& \Pi [(2J'_{r0} + 1)(2J_{r0} + 1)]^{1/2} \Pi \Delta(J_{r0} j J_{r+1,0}) \Pi \Delta(J'_{r0} j J'_{r+1,0}) \times \\
& \Delta(J_{n-1,0} j J_n) \Delta(J'_{n-1,0} j J_n) \times \\
& N(J_{n-1,0} J_n) \Pi N(J_{r0} J_{r+1,0}) \Pi N(J_{r1} J_{r+1,1}) \Pi [(2J_{r0} + 1)(2J_{r1} + 1)]^{1/2} \times \\
& \Pi \Delta(J_{r0} j J_{r+1,0}) \Pi \Delta(J_{r1} j J_{r+1,1}) \Delta(J_{n-1,0} j J_n) \Delta(J_{n-1,1} j J_n)
\end{aligned} \tag{36}$$

In (36), all terms with the various J_{r0} are squared and may be absorbed in

the rational function which multiplies the following expression

$$\begin{aligned}
& N(J'_{n-1,0}J_n)\Pi N(J'_{r0}J'_{r+1,0})\Pi N(J_{r1}J_{r+1,1})\times \\
& \Pi[(2J'_{r0}+1)(2J_{r1}+1)]^{1/2}\Pi\Delta(J'_{r0}jJ'_{r+1,0})\Pi\Delta(J_{r1}jJ_{r+1,1})\times \\
& \Delta(J'_{n-1,0}jJ_n)\Delta(J_{n-1,1}jJ_n)
\end{aligned} \tag{37}$$

The expression (37) is identical with the corresponding c.f.p. of the state with J'_{r0} multiplied by a rational function. Hence, also the c.f.p. of the orthogonalized state are square roots of rational functions and have the form given by (33). All states with the same value of J_n have c.f.p. given by (33). They are different by a common factor to all c.f.p. of each state which is a square root of a rational function. They are orthogonal to each other due to the different rational functions multiplying the expression (33) to obtain the full c.f.p. Some of these rational functions may vanish. The states orthogonalized in this way are not normalized and must be explicitly normalized.

We turn now to the form (33) of c.f.p. of states constructed consecutively from principal parents and antisymmetrized. Obviously, it holds for the case $n=3$. To demonstrate that it is correct also for c.f.p. with $n > 3$ we use the recursion relation (31) for induction by n . We show that if (31) is the correct form for a given $n-1$, it holds also for n . According to the assumption, every term in the summation in (31) is given by a rational function multiplying the expression

$$\begin{aligned}
& N(J_{n-2,0}J_{n-1,0})\Pi N(J_{r0}J_{r+1,0})\Pi N(J_{r1}J_{r+1,1})\Pi[(2J_{r0}+1)(2J_{r1}+1)]^{1/2}\times \\
& N(J'_{n-2,0}J'_{n-1,0})\Pi N(J'_{r0}J'_{r+1,0})\Pi N(J_{r1}J_{r+1,1})\Pi[(2J'_{r0}+1)(2J_{r1}+1)]^{1/2}\times \\
& \Pi\Delta(J_{r0}jJ_{r+1,0})\Pi\Delta(J_{r1}jJ_{r+1,1})\Delta(J_{n-2,0}jJ_{n-1,0})\Delta(J_{n-2,1}jJ_{n-1,0})\times \\
& \Pi\Delta(J'_{r0}jJ'_{r+1,0})\Pi\Delta(J_{r1}jJ_{r+1,1})\Delta(J'_{n-2,0}jJ'_{n-1,0})\Delta(J_{n-2,1}jJ'_{n-1,0})\times \\
& \{J_{n-2,1}jJ'_{n-1,0}/J_njJ_{n-1,0}\}
\end{aligned} \tag{38}$$

This expression may be simplified by noting that there are several identi-

cal factors which may be absorbed in the rational function. Thus, we obtain instead of (38) the form

$$\begin{aligned}
& \Pi N(J_{r1}J_{r+1,1})^2 \Pi(2J_{r1}+1) \Pi \Delta(J_{r1}jJ_{r+1,1})^2 \times \\
& \Pi N(J_{r0}J_{r+1,0}) \Pi N(J'_{r0}J'_{r+1,0}) \Pi[(2J_{r0}+1)(2J'_{r0}+1)]^{1/2} \times \\
& \Pi \Delta(J_{r0}jJ_{r+1,0}) \Pi \Delta(J'_{r0}jJ'_{r+1,0}) \{J_{n-2,1}jJ'_{n-1,0}/J_njJ_{n-1,0}\} \times \\
& \Delta(J_{n-2,1}jJ_{n-1,0}) \Delta(J_{n-2,1}jJ'_{n-1,0})
\end{aligned} \tag{39}$$

Whereas in the products in (38) r goes up to $n-3$, in (39) it goes up to $n-2$. Using the formula (7) for the $6j$ -symbol, the product

$$\Delta(J_{n-2,1}jJ'_{n-1,0}) \Delta(J_{n-2,1}jJ_{n-1,0}) \{J_{n-2,1}jJ'_{n-1,0}/J_njJ_{n-1,0}\} \tag{40}$$

in (39) becomes equal to a rational function multiplied by

$$\Delta(J_{n-2,1}jJ_{n-1,0})^2 \Delta(J_{n-2,1}jJ'_{n-1,0})^2 \Delta(J_{n-1,0}jJ_n) \Delta(J'_{n-1,0}jJ_n) \tag{41}$$

Thus the $J_{n-2,1}$ angular momentum appears as an argument only of rational functions and the summation over it can be directly carried out for all terms. The result of this summation is a rational function multiplied by

$$\begin{aligned}
& \Pi N(J_{r0}J_{r+1,0}) \Pi N(J'_{r0}J'_{r+1,0}) \Pi[(2J_{r0}+1)(2J'_{r0}+1)]^{1/2} \times \\
& \Pi \Delta(J_{r0}jJ_{r+1,0}) \Pi \Delta(J'_{r0}jJ'_{r+1,0}) \Delta(J_{n-1,0}jJ_n) \Delta(J'_{n-1,0}jJ_n)
\end{aligned} \tag{42}$$

According to (31), this expression should be multiplied by $N(J_{n-1,0}J_n)$ and is then identical with (33).

Thus, if the form (33) is correct for $n-1$ it is correct also for n . Since it is correct for $n=3$, it holds for any value of n . From (33) follows that the squares of c.f.p. obtained from principal parents and antisymmetrization are rational functions. According to (3), this is sufficient to guarantee that if in every j^r configuration, $r \leq n$, there is only one state with a given value of J_r , the diagonal matrix elements have the property considered in this paper. This special case was discussed in the Introduction in a simpler way. Relation (3) becomes more complicated if there are several states with the same value of

J . Unlike the situation in the simple $n=3$ case, using relation (3) for diagonal matrix elements in the j^n configuration may well lead to non-diagonal matrix elements in j^r configurations with $r \leq n$. Non-diagonal matrix elements are not given by squares of $n-1 \rightarrow n$ c.f.p. There is, however, a way to make use of the structure (33) of these c.f.p. to show that the states constructed in this way have the property considered in this paper.

Any antisymmetric wave function in the j^n configuration may be expanded in $n-2 \rightarrow n$ c.f.p. which are discussed in detail in ref.[2]. They are defined by

$$\psi(j^n aJ) = \Sigma[j^{n-2}(bJ_1)j^2(J')J]\{j^n aJ\}\psi(j^{n-2}(bJ_1)j_{n,n-1}^2(J')aJ) \quad (43)$$

Such an expansion is always possible since the wave functions on the r.h.s. of (43) form a complete basis for the space of j^n states with given J , which are antisymmetric in particles 1 to $n-2$ and in particles $n, n-1$. Fully antisymmetric j^n wave functions with given J are included in this space and hence, may be expanded as in (43). The calculation of two-body matrix elements of a scalar operator, like the mutual interaction between particles, is very simple if such c.f.p. are used. Since these states are fully antisymmetric, it is possible to calculate matrix elements of one term in the interaction, $V_{n,n-1}$ say, and multiply the result by the number of such terms, $n(n-1)/2$. Thus, using the expansion (43), we obtain the result

$$\begin{aligned} < j^n aJ | \Sigma V_{ik} | j^n aJ > = n(n-1) < j^n aJ | V_{n,n-1} | j^n aJ > / 2 = \\ & [n(n-1)/2] \Sigma[j^{n-2}(bJ_1)j^2(J')J]\{j^n aJ\}[j^{n-2}(b''J_1'')j^2(J'')J]\{j^n aJ\} \times \\ & < j^{n-2}(bJ_1)j^2(J')J | V_{n,n-1} | j^{n-2}(b''J_1'')j^2(J'')J > \end{aligned} \quad (44)$$

Integration of (44) over coordinates 1 to $n-2$ yields zero unless $J_1 = J_1''$ and $b'' \equiv b$. Since $V_{n,n-1}$ is a scalar operator, its matrix element on the r.h.s. of (44) vanishes if J'' is not equal to J' . If they are equal, the two-body matrix element in (44) is equal to $< j^2 J' | V | j^2 J' >$. Hence, the matrix

element is given by

$$\begin{aligned} & \langle j^n aJ | \Sigma V_{ik} | j^n aJ \rangle = \\ & [n(n-1)/2] \Sigma_{bJ_1 J_1'} [j^{n-2}(bJ_1)j^2(J')J] \{j^n aJ\}^2 \langle j^2 J' | V j^2 J' \rangle \end{aligned} \quad (45)$$

Hence, using the $n-2 \rightarrow n$ c.f.p. in (43), diagonal matrix elements of a two-body interaction are given by the linear combination (45). In it, the coefficients of the two-body energies are squares of the c.f.p. multiplied by $n(n-1)/2$. The expectation value in the state $|j^n aJ\rangle$ of a two-body interaction is a linear combination of two-body energies with coefficients that are positive rational functions if the squares of these c.f.p. are rational.

To find out which states have $n-2 \rightarrow n$ c.f.p. whose squares are rational functions, it is convenient to express them in terms of $n-1 \rightarrow n$ c.f.p. This expression is given by (26.44) in ref.[2] as

$$\begin{aligned} & [j^{n-2}(a_{n-2,0}J_{n-2,0})j^2(J')J_n] \{j^n aJ_n\} = \\ & \Sigma [j^{n-2}(a_{n-2,0}J_{n-2,0})jJ_{n-1,1}] \{j^{n-1}a_{n-1,1}J_{n-1,1}\} \times \\ & [J^{n-1}(a_{n-1,1}J_{n-1,1})jJ_n] \{j^n aJ_n\} \times \\ & [(2J_{n-1,1} + 1)(2J' + 1)]^{1/2} \{J_{n-2,0}jJ_{n-1,1}/jJ_n J'\} \end{aligned} \quad (46)$$

The form (33) was proved above to be valid for c.f.p. of certain states constructed from principal parents. Using such states in (46), the form of their $n-1 \rightarrow n$ c.f.p. may be taken from (33). Thus, the $n-2 \rightarrow n-1$ c.f.p. $[j^{n-2}(a_{n-2,0}J_{n-2,0})jJ_{n-1,1}] \{j^{n-1}[a_{n-2,1}, J_{n-2,1}]J_{n-1,1}\}$ is expressed as a rational function multiplying

$$\begin{aligned} & N(J_{n-2,1}J_{n-1,1})\Pi N(J_{r1}J_{r+1,1})\Pi N(J_{r0}J_{r+1,0})\Pi[(2J_{r1} + 1)(2J_{r0} + 1)]^{1/2} \times \\ & \Pi\Delta(J_{r1}jJ_{r+1,1})\Pi\Delta(J_{r0}jJ_{r+1,0})\Delta(J_{n-2,1}jJ_{n-1,1})\Delta(J_{n-2,0}jJ_{n-1,1}) \end{aligned} \quad (47)$$

The $n-1 \rightarrow n$ c.f.p. $[j^{n-1}([a_{n-2,1}, J_{n-2,1}]J_{n-1,1})jJ_n] \{j^n[J'_{n-1,0}]J_n\}$ in (46) can be similarly expressed, as a rational function multiplying the following

expression which is written in more detail than in (33)

$$\begin{aligned}
& N(J'_{n-1,0}J_n)\Pi N(J'_{r0}J'_{r+1,0})\Pi N(J_{r1}J_{r+1,1})N(J_{n-2,1}J_{n-1,1})\times \\
& \Pi[(2J'_{r0}+1)(2J_{r1}+1)]^{1/2}[(2J'_{n-1,0}+1)(2J_{n-1,1}+1)]^{1/2}\times \\
& \Pi\Delta(J'_{r0}jJ'_{r+1,0})\Pi\Delta(J_{r1}jJ_{r+1,1})\Delta(J_{n-2,1}jJ_{n-1,1})\times \\
& \Delta(J'_{n-1,0}jJ_n)\Delta(J_{n-1,1}jJ_n)
\end{aligned} \tag{48}$$

Substituting these expressions into (46) yields for each term in the summation a rational function multiplied by

$$\begin{aligned}
& N(J'_{n-1,0}J_n)N(J_{n-2,1}J_{n-1,1})^2 \times \Pi N(J_{r1}J_{r+1,1})^2 \Pi N(J_{r0}J_{r+1,0}) \\
& \Pi N(J'_{r0}J'_{r+1,0})\Pi[(2J_{r0}+1)(2J'_{r0}+1)]^{1/2}\Pi(2J_{r1}+1)\times \\
& [(2J'_{n-1,0}+1)(2J'+1)]^{1/2}\Pi\Delta(J_{r0}jJ_{r+1,0})\Pi\Delta(J'_{r0}jJ'_{r+1,0})\times \\
& \Pi\Delta(J_{r1}jJ_{r+1,1})^2\Delta(J_{n-2,0}jJ_{n-1,1})\Delta(J_{n-2,1}jJ_{n-1,1})^2\Pi\Delta(J_{r0}jJ_{r+1,0})\times \\
& \Delta(J'_{n-1,0}jJ_n)\Delta(J_{n-1,1}jJ_n)(2J_{n-1,1}+1)\{J_{n-2,0}jJ_{n-1,1}/jJ_nJ'\}
\end{aligned} \tag{49}$$

Several terms in (49) are rational functions. Using the form (7) of $6j$ -symbols, the following product

$$\Delta(J_{n-2,0}jJ_{n-1,1})\Delta(J'_{n-1,0}jJ_n)\Delta(J_{n-1,1}jJ_n)\{J_{n-2,0}jJ_{n-1,1}/jJ_nJ'\} \tag{50}$$

becomes equal to a rational function multiplying

$$\begin{aligned}
& \Delta(J'_{n-1,0}jJ_n)\Delta(J_{n-1,1}jJ_n)^2\Delta(J_{n-2,0}jJ_{n-1,1})^2\times \\
& \Delta(jjJ')\Delta(J_{n-2,0}J'J_n)
\end{aligned} \tag{51}$$

Thus, (49) may be expressed as a rational function multiplied by

$$\begin{aligned}
& N(J'_{n-1,0}J_n)\Pi N(J_{r0}J_{r+1,0})\Pi N(J'_{r0}J'_{r+1,0})\Pi[(2J_{r0}+1)(2J'_{r0}+1)]^{1/2} \\
& [(2J'_{n-1,0}+1)(2J'+1)]^{1/2}\Pi\Delta(J'_{r0}jJ'_{r+1,0})\Pi\Delta(J_{r1}jJ_{r+1,1}) \\
& \Delta(J'_{n-1,0}jJ_n)\Delta(jjJ')\Delta(J_{n-2,0}J'J_n)
\end{aligned} \tag{52}$$

The summation in (46) is over $J_{n-1,1}$ which does not appear in (52) and hence, it appears only in the rational function multiplying it. Therefore, the

summation over it in (46) can be simply carried out yielding another rational function multiplying the expression (52).

Thus, the $n-2 \rightarrow n$ c.f.p. $[j^{n-2}(a_{n-2,0}J_{n-2,0})j^2(J')J_n|\}j^n a J_n]$ is the square root of a rational function and its square is a rational function. According to (45), the expectation value of a state with these c.f.p. of a two-body interaction is a linear combination of two-body energies with rational coefficients.

The $n-2 \rightarrow n$ c.f.p. obtained above, may be used to derive expressions of non-diagonal matrix elements of a two-body interaction. Instead of (45), such non-diagonal elements are defined by

$$\begin{aligned} < j^n a J | \Delta V_{ik} | j^n a' J > = \\ & [n(n-1)/2] \Sigma [j^{n-2}(bJ_1)j^2(J')J|\}j^n a J] \times \\ & [j^{n-2}(bJ_1)j^2(J')J|\}j^n a' J] < j^2 J' | V j^2 J' > \end{aligned} \quad (53)$$

Only non-vanishing terms are written on the r.h.s. of (53). The expression (52) is taken to be the non-rational factor in the c.f.p. of the state a . The c.f.p. of the state a' is the same expression in which J'_{r0} is replaced by J_{r0} . All angular momenta J_{r0} which define the $|j^{n-2}bJ_{n-2,0}>$, as well as J' , appear in the same expressions in both c.f.p. and hence, contribute in (53) only rational functions. Each term in the summation in (53) is thus equal to a rational function multiplying the expression

$$\begin{aligned} & N(J'_{n-1,0}J'_n)N(J_{n-1,0}J_n)\Pi N(J'_{r0}J'_{r+1,0})\Pi N(J_{r0}J_{r+1,0}) \times \\ & \Pi[(2J'_{r0}+1)(2J_{r0}+1)]^{1/2}\Pi\Delta(J'_{r0}jJ'_{r+1,0})\Pi\Delta(J_{r0}jJ_{r+1,0}) \times \\ & \Delta(J'_{n-1,0}jJ_n)\Delta(J_{n-1,0}jJ_n)V(J') \end{aligned} \quad (54)$$

The factor of $V(J')$ in (54) is a common factor of all terms in the summation in (53). Hence, the non-diagonal matrix element (53) is a linear combination of two-body energies whose coefficients are rational functions multiplied by a common, J' independent factor. The square of this factor as seen in (54) is a rational function of the angular momenta involved.

4 The seniority scheme

The seniority quantum number v measures in some sense the amount of pairing of particle pairs in $J=0$ states. In the seniority scheme, states are eigenstates of the pairing interaction defined by

$$\langle j^2 JM \mid q_{12} \mid j^2 JM \rangle = \delta_{J0} \delta_{M0} (2j + 1) \quad (55)$$

The eigenvalues of this interaction in the j^n configuration are equal to

$$(n - v)(2j + 3 - n - v)/2 \quad (56)$$

Thus, loosely speaking, v is the number of unpaired particles. There is a very large class of interactions, not just the pairing interaction, which are diagonal in the seniority scheme. Also for general interactions, the seniority scheme is a convenient one to use. Here, this scheme will be discussed only in relation to the theme of the paper. The aim is to see which of the states of this scheme have diagonal matrix elements which are linear combinations of two-body energies with rational coefficients.

According to (56), states with seniority v in the j^v configuration have no $J=0$ pairing. States with the same seniority in higher configurations are obtained by starting from those states, adding $(n - v)/2$ pairs with $J=0$ and antisymmetrizing. Adding one particle to a state with seniority v and antisymmetrizing, the state obtained has seniority $v+1$ or $v-1$. It may also be a linear combination of states with both seniorities. The state with no j -particles has seniority $v=0$ whereas the one particle state, with $J=j$, has $v=1$. States with two particles have states with $J=0$, $v=0$ or $J > 0$, *even* and seniority $v=2$.

Consider a state in the j^n configuration, $n < 2j$ *even* with $J=0$ and $v=0$. Adding one particle to this principal parent, a state with $J=j$ and $v=1$ is obtained. A state, with $n < 2j+1$ *odd*, with $J = j$, $v=1$, is a principal parent of a $J=0$, $v=0$ state. Hence, such states have the property considered in this paper. The expectation values of any two-body interaction of states

with lowest seniorities, $v=0$ or $v=1$, are given explicitly by (30.1) ref.[2] or (20.13) ref.[5] as

$$an(n-1)/2 + [n/2]b \quad (57)$$

where $[n]$ is the largest integer not exceeding n . In (57), a and b are linear combinations of the $V(J)$ which appear only in two linear combinations

$$V_0 = V(J=0) \quad \text{and} \quad \bar{V}_2 = \Sigma(2J+1)V(J)/\Sigma(2J+1) \quad J > 0, \text{even}$$

The coefficients a and b are given by these combinations as

$$a = \{(2j+2)\bar{V}_2 - V_0\}/(2j+1) \quad b = (2j+2)(V_0 - \bar{V}_2)/(2j+1) \quad (58)$$

In the case of an interaction which is diagonal in the seniority scheme, the expressions (57) are the eigenvalues of these states. Unlike the cases of $j < 7/2$, there are no closed formulae for diagonal matrix elements of states with higher values of v . Similar expressions exist only for the average values of diagonal matrix elements of all states with given v in the j^n configuration $v = n, n-2, \dots$. It will be shown, however, that states of the seniority scheme may be constructed from principal parents, antisymmetrizations and orthogonalization if necessary. As shown in the preceding section, this guarantees that they have the property considered here. Diagonal matrix elements of a two-body interaction of states in the seniority scheme, constructed as explained above, are linear combinations of two-body energies with rational coefficients.

In the j^2 configuration, the $J=0$ state has seniority $v=0$ as mentioned above. The other antisymmetric states have even values of $J > 0$ and seniority $v=2$. As mentioned above, adding a j -particle to a principal parent with $J=0$, $v=0$, the state j^n $J=j$ with seniority $v=1$ is obtained. Since $n-v$ is always an even integer, adding a j -particle to a state with seniority v in the j^n configuration (and antisymmetrizing) yields a state of the j^{n+1} configuration with seniority $v+1$ or $v-1$ or both. The added particle either completes a

$J=0$ pair in which case the seniority is reduced or destroys a $J=0$ pair which raises the seniority.

If a j -particle is added to a principal parent with $J_0 > 0$, $v=2$, the state obtained has $v=3$ if $J \neq j$. If a state with $J = j$ is constructed, it is a linear combination of states with seniorities $v=1$ and $v=3$, if there is a state with $J = j$, $v=3$ in the j^3 configuration. Such a linear combination will have a non-vanishing c.f.p. $[j^2(v=0, J=0)J=j|j^3J=j]$. The $J = j$, $v=1$ state obtained earlier, may be projected out by adding it with a coefficient that was defined by (13). The remaining state will be a pure $J = j$, $v=3$ state, orthogonal to the $J = j$, $v=1$ state. This procedure was described in detail in the preceding section. If there are several $J = j$ states in the j^3 configuration, they may be constructed from several principal parents. They should then be orthogonalized to the $J = j$, $v=1$ state and among themselves.

This procedure may be carried out in any j^n configuration. If n is even, the steps described above may be taken. States with seniorities $v=4$ to which a j -particle is added, may be used to construct $v=5$ states or linear combinations of $v=3$ and $v=5$ states. In that case, the components with $v=3$ may be projected out by using the $v=3$ states constructed in the preceding step. At the end, states with seniority $v = n$ may be used to construct states with seniorities $v = n+1$ in the j^{n+1} configuration.

If n is odd, the state with $J = j$, $v=1$ may be used to construct a state with $J=0$ which has seniority $v=0$, there are no states with $J=0$ and $v=2$. States with seniority $v=2$ may be constructed from the same $J = j$, $v=1$ state by coupling an additional j -particle to obtain a state with $2j > J > 0$, *even*. From states with $v=3$, states with $v=4$ may be obtained. If a linear combination of $v=2$ and $v=4$ is obtained, the $v=2$ components may be projected out by using $v=2$ states obtained earlier. This procedure may be further carried out until all states in the j^{n+1} configuration are constructed. They all have well defined seniorities and their diagonal matrix elements have the property described in this paper.

An example of the procedure described above is offered in ref.[1]. The authors calculated the eigenvalues of $J=2$ and $J=4$ states with seniority $v=4$ in the $(7/2)^4$ configuration. They found that the coefficients of two-body energies in both cases are rational numbers. In that configuration, there are also $J=2$ and $J=4$ states with seniority $v=2$. The latter may be obtained from the state in where a $7/2$ particle is coupled to the principal parent with $J=7/2$ in the $(7/2)^3$ configuration. There is only one such state in that configuration and hence, it has seniority $v=1$. The $J=2$ and $J=4$ states obtained from it have seniority $v=2$ since the only state with $v=0$ has $J=0$. The other $J=2,4$ states must be obtained from another principal parent. In that one, a $7/2$ particle should be coupled with any $v=3$ state in the $(7/2)^3$ configuration. From the resulting states, the $v=2$ components should be projected out as described above. Hence, the $v=4$ states are examples of states which have the property considered in the present paper.

A more complicated situation occurs in the $(9/2)^4$ configuration considered in ref.[1]. In that case there are 3 states with $J=4$ and 3 states with $J=6$. One of the $J=4$ (and $J=6$) states has seniority $v=2$ and may be constructed from a state where a $9/2$ particle is coupled to the $J=9/2$, $v=1$ state. One of the remaining $J=4$ states may be constructed from a principal parent with $J=9/2$, $v=3$ coupled to a $9/2$ particle. The resulting state, $|a\rangle$, is a linear combination of $J=4$, $v=4$ and $J=4$, $v=2$ states. Another independent $J=4$ state, $|b\rangle$, may be constructed from another principal parent with $v=3$ and is also a combination of $v=4$ and $v=2$ states. The $J=4$, $v=4$ states which are discussed in ref.[1] may be obtained as follows. First, the state $|b\rangle$ is orthogonalized to state $|a\rangle$ as indicated by vanishing of the c.f.p. $[(9/2)^3(J_0 = 9/2, v = 3)(9/2)J = 4]\{(9/2)^4 b J = 4\}$ of the orthogonalized $|b\rangle$ state. Then, from both states the $J=4$, $v=2$ components are projected out leaving them as pure $v=4$ states. In each resulting state the c.f.p. corresponding to that of the principal parent of the $v=2$ state, $[(9/2)^3(J_0 = 9/2, v = 1)(9/2)J = 4]\{(9/2)^4 J = 4\}$, vanishes. It follows that these $v=4$ states have the property considered in this paper. The same arguments apply also to the $J=6$, $v=4$ states. Their diagonal matrix elements are linear combinations of two-body energies with rational coefficients.

5 Summary

In the present paper, matrix elements of a two-body interaction between states of j^n configurations were considered. The two-body interaction is taken to be a perturbation and hence, only the sub-matrix of the Hamiltonian, defined by states of the j^n configuration, is considered. The matrix elements are functions of two-body energies $V(J) = \langle j^2 JM | V | j^2 JM \rangle$. There are states whose expectation values (diagonal matrix elements) are linear combinations of the $V(J)$ with positive rational coefficients. In this paper, it was shown that this is always the case for states which are the only ones with given J in the j^n configuration considered. Such states are eigenstates of the sub-matrix considered.

When there are several states with the same value of J , the situation is more complicated. It was shown above that it is possible to construct in such cases, a complete basis of states which have this simple feature. The diagonal matrix elements of these states are linear combinations of the $V(J)$ whose coefficients are positive rational numbers. The construction of these states was carried out by using coefficients of fractional parentage (c.f.p.). These are obtained by starting from states of the j^3 configuration which serve as principal parents. This procedure is repeated by adding more particles until the j^n configuration is reached. In every step, orthogonalization (and normalization) of the states thus obtained, is performed. It was shown that the squares of the c.f.p. are rational functions of the angular momenta used in the construction and hence, are independent of the particular two-body interaction.

It was shown that states of the seniority scheme form bases of this kind. If the interaction is diagonal in the seniority scheme, non-diagonal matrix elements between states with different seniorities vanish. This, however, is not true for the most general two-body interaction. In any case, there are usually non-vanishing non-diagonal matrix elements between states with the same seniority v and the same spin J . Irrespective of the nature of the basis

states for a given value of J , they define a sub-matrix of the Hamiltonian. This sub-matrix should be diagonalized in order to find the eigenstates and corresponding eigenvalues.

The eigenstates are linear combinations of the basis states but the coefficients strongly depend on the eigenvalues. The latter are functions of the $V(J)$ but are usually far from linear functions. The eigenvalues are the roots of the secular equation whose degree is limited only by the order of the sub-matrix. The eigenstates may be expanded in terms of c.f.p. but unlike the basis states constructed in this paper, their c.f.p. are functions of the $V(J)$.

It is convenient to use the basis states to set up the sub-matrix to be diagonalized. In addition to the rather simple structure of diagonal elements, also the non-diagonal elements have a simple structure. Each is a linear combination of the $V(J)$ with rational coefficients multiplied by a factor which is the square root of a rational function

From the discussion above, a simple conclusion follows that states with the rational property are the exception rather than the rule. In any given j^n configuration, states which are the only ones with a given value of J are rather rare. They may be found among the states with the highest values of J (certainly the state $J = nj - n(n-1)/2$) as well as among those with the lowest J values.

Still, there may be special cases like those mentioned in ref.[1]. In the case of $j = 7/2$, the non-diagonal matrix element of any two body interaction between the $J=2$ states with $v=2$ and $v=4$ vanishes. The same is true also for the $v=2$ and $v=4$ states with $J=4$. This follows from general properties of the seniority scheme. More surprising is the fact that in the $(9/2)^4$ configuration, there is a $v=4$, $J=4$ state whose matrix element of any two-body interaction, vanishes not only with the $v=2$, $J=4$ state but also with the orthogonal $v=4$, $J=4$ state. Its eigenvalue is a linear combination of two-body interactions with positive rational coefficients as given in ref.[1]. The same situation

occurs also for the corresponding $J=6$ states.

References

- [1] L.Zamick and P.Van Isacker, Phys. Rev. **78**, 044327 (2008).
- [2] A.de Shalit and I.Talmi, Nuclear Shell Theory, Academic Press (1963);
Reprinted by Dover Publications (2003).
- [3] R.F.Bacher and S.Goudsmit, Phys. Rev. **46** (1934) 948.
- [4] G.Racah, Phys. Rev. **63** (1943) 367.
- [5] I.Talmi, Simple Models of Complex Nuclei, Harwood Academic Publishers (1993).
- [6] G.Racah, Phys. Rev. **62** (1942) 438.